

Probiotics for Metabolic Health

Today, a record number of patients worldwide suffer from metabolic disorders, including obesity, non-alcoholic fatty liver disease (NAFLD), type 2 diabetes mellitus (T2DM) and cardiovascular disease (CVD). As poor diets, lack of exercise, and other stressors continue to negatively impact millions of people around the globe, we must look for new ways to improve metabolic health, delay disease progression, and foster a better quality of life where possible.

The microbiota and metabolic disorders

Currently there is no strict definition for metabolic health. Often it is referred to as the absence of metabolic disorders. Rising levels of blood sugar, triglycerides, low-density lipoprotein LDL cholesterol, blood pressure, and waist circumference are generally associated with loss of metabolic health and an increased risk for metabolic diseases ranging from mild insulin resistance all the way to prediabetes and type 2 diabetes mellitus and cardiovascular disease². Both insulin resistance and systemic low-grade inflammation seem to be at the core of metabolic disorders³⁻⁵.

Insulin resistance is a condition in which cells are less sensitive to the action of insulin. Thus, the pancreas needs to produce more insulin to lower blood glucose levels. Over time, the pancreas may no longer be able to cope with the high demand, and prediabetes can progress to diabetes^{3,4}.

Recent research has indicated that the gut microbiota plays an important role in metabolic health⁶. Disturbance of gut microbiota due to e.g. a typical western lifestyle leads to changes in the serum lipopolysaccharides (LPS), short-chain fatty acids (SCFAs) and bile acid, resulting in systemic low-grade inflammation and insulin resistance⁷⁻⁹ [figure 1].

Is there a role for probiotics?

Given the role of the gut microbiota on metabolic disorders, targeted probiotic formulations may be clinically relevant for optimizing metabolic health, influencing of insulin resistance and low-grade inflammation associated with early- and late-stage low-grade metabolic disorders, specifically T2DM. Recent literature has supported the efficacy of probiotics for improving a range of metabolic markers, including HOMA-IR, a measure of insulin resistance, and serum LPS, a measure of gut permeability and a trigger of inflammatory responses¹⁰⁻¹⁴.

Ecologic® Barrier for metabolic health

The probiotic formulation Ecologic® Barrier has been developed by Winlove Probiotics already in 2012. Ecologic® Barrier is a multispecies probiotic developed to optimize Barrier function and to reduce systemic low-grade inflammation¹⁵, making it a suitable choice for research and application in insulin resistance and metabolic health. The probiotic contains the following bacterial strains: *B. bifidum* W23, *B. lactis* W51, *B. lactis* W52, *L. acidophilus* W37, *L. brevis* W63, *L. casei* W56, *L. salivarius* W24, *Lc. lactis* W19, *Lc. Lactis* W58 with a viable cell count of 2.5×10^9 cfu/gram.




FIGURE 1: Crosstalk between the gut microbiota and the host's system in terms of inflammation and metabolism. The gut microbiota, through a range of molecular interactions, contributes to insulin resistance, systemic low-grade inflammation, and fat deposition and therefore, indirectly participates in the onset and progress of metabolic disorders.

The probiotic strains were selected based on the following criteria: in vitro strengthening of the epithelial barrier, inhibition of mast cell activation, inhibition of pro-inflammatory cytokines and decreasing lipopolysaccharide load.

Clinical evidence

Since 2017, a series of randomized, double-blind, placebo-controlled studies have established the value of Ecologic® Barrier for the improvement of metabolic health.

Ecologic® Barrier has been tested in a double-blind, placebo-controlled, randomized study, performed by the Warwick University, UK and King Saud University, Saudi Arabia^{10,11}. Ninety-six adult T2DM patients (treatment-naïve and without comorbidities) were randomized to receive 2 grams of Ecologic® Barrier or placebo twice daily [1.0×10^{10} cfu/day] for 6 months. In the probiotic group Ecologic® Barrier significantly reduced HOMA-IR levels after 3 months and 6 months, which did not occur in the placebo group (figure 2). In line with this, a significant decrease in fasting glucose and fasting insulin was observed in the probiotic group. In addition, Ecologic® Barrier intake reduced circulating endotoxin levels [LPS], a trigger of inflammation and a marker for Barrier function, and improved inflammation markers such as CRP (figure 3), TNF- α , IL-6. The positive effect of Ecologic® Barrier on the gut Barrier function was also observed in a study

performed by researchers from the Medical University of Graz, Austria¹⁴. This randomized, double-blind, placebo-controlled pilot study investigated the effect of Ecologic® Barrier combined with a prebiotic on glucose metabolism, gut microbiota, and gut permeability in obese patients with T2DM. For this study, twenty-six treatment-experienced obese T2DM patients were randomized to daily receive 6 grams of Ecologic® Barrier [1.5×10^{10} cfu/day] plus a prebiotic or a placebo for 6 months. After 3 months patients in the placebo group showed a worsened gut permeability [increase in serum zonuline] which was not observed in the Ecologic® Barrier plus prebiotic group.

Another double-blind, placebo-controlled randomized study performed by the University of Medical Sciences in Poznan, Poland studied the effects of Ecologic® Barrier on the metabolic health of obese postmenopausal women¹². Eighty-one obese postmenopausal women were randomly assigned to receive placebo, a low dose of Ecologic® Barrier [LD] [2.5×10^9 cfu/day], or a high dose of Ecologic® Barrier [HD] [1×10^{10} cfu/day] divided into two equal doses for 12 weeks. Both LD and HD Ecologic® Barrier intake resulted in significantly reduced HOMA-IR levels compared to baseline, which was not observed in the placebo group. A dose-response effect was observed as a significant larger reduction of HOMA-IR occurred in the




FIGURE 2: HOMA-IR levels (Median [range]) before and after 3 months and 6 months supplementation with Ecologic® Barrier.

* Significant decrease, $p < 0.05$




FIGURE 3: C-reactive protein (CRP) (Median [range]) before and after 3 months and 6 months supplementation with Ecologic® Barrier.

* Significant decrease, $p < 0.05$

HD group [figure 4]. Moreover, Ecologic® Barrier improved circulating endotoxin [figure 5]. A second publication of the same clinical trial showed that inflammation makers such as TNF- α , IL-6 and functional and biochemical markers of vascular dysfunction such as blood pressure improved as well¹³.

Future clinical applications

Taken together, these studies establish a strong rationale for Ecologic® Barrier’s impact on key metabolic markers, both biochemical and functional. Several other probiotics have been studied for their effect on insulin resistance and type 2 diabetes. Unlike many other formulations, Ecologic® Barrier has been investigated in several independent studies showing a clear positive effect on HOMA-IR, which was dose and duration-dependent. This positive effect was also observed on markers of inflammation [e.g. CRP,

TNF- α and IL-6], intestinal Barrier function [zonulin and LPS] and vascular dysfunction [e.g. blood pressure]. Overall, this indicates that Ecologic® BARRIER can lower both insulin resistance and systemic low-grade inflammation, which are the hallmark symptoms in metabolic disorders such as prediabetes and T2DM. This may help to either postpone, prevent, or lower the need for medication in prediabetic patients, and might help to lower medication class or dosage in diabetics. In conclusion, these studies show that there is a role to play for Ecologic® Barrier in improving overall metabolic health, ideally when combined with a comprehensive lifestyle, wellness, and medical plan.




FIGURE 4: HOMA-IR levels (Mean \pm SD) before and after 12 weeks of low or high dose supplementation with Ecologic® Barrier.

* Significant decrease, $p < 0.05$




FIGURE 5: LPS levels (Mean \pm SD) before and after 12 weeks of low or high dose supplementation with Ecologic® Barrier.

* Significant decrease, $p < 0.05$

We encourage you to learn more about the Ecologic® Barrier formulation and its wide-ranging health benefits. Reach out to the team at Winclove Probiotics.

References

- 1 Bennett JE, Stevens GA, Mathers CD, et al. NCD Countdown 2030: worldwide trends in non-communicable disease mortality and progress towards Sustainable Development Goal target 3.4. *The Lancet* 2018; 392: 1072–88.
- 2 Saklayen MG. The Global Epidemic of the Metabolic Syndrome. *Curr Hypertens Rep* 2018; 20. DOI:10.1007/s11906-018-0812-z.
- 3 Ighbariya A, Weiss R. Insulin Resistance, Prediabetes, Metabolic Syndrome: What Should Every Pediatrician Know? *J Clin Res Pediatr Endocrinol* 2018; : 49–57.
- 4 Petersen MC, Shulman GI. Mechanisms of Insulin Action and Insulin Resistance. *Physiol Rev* 2018; 98: 2133–223.
- 5 Kumari R, Kumar S, Kant R. An update on metabolic syndrome: Metabolic risk markers and adipokines in the development of metabolic syndrome. *Diabetes Metab Syndr Clin Res Rev* 2019; 13: 2409–17.
- 6 Parekh PJ, Balart LA, Johnson DA. The Influence of the Gut Microbiome on Obesity, Metabolic Syndrome and Gastrointestinal Disease. *Clin Transl Gastroenterol* 2015; 6: e91.
- 7 Janssen AWF, Kersten S. The role of the gut microbiota in metabolic health. *FASEB J Off Publ Fed Am Soc Exp Biol* 2015; 29: 3111–23.
- 8 Dabke K, Hendrick G, Devkota S. The gut microbiome and metabolic syndrome. *J Clin Invest* 2019; 129: 4050–7.
- 9 Jayashree B, Bibin YS, Prabhu D, et al. Increased circulatory levels of lipopolysaccharide (LPS) and zonulin signify novel biomarkers of proinflammation in patients with type 2 diabetes. *Mol Cell Biochem* 2014; 388: 203–10.
- 10 Sabico S, Al-Mashharawi A, Al-Daghri NM, et al. Effects of a multi-strain probiotic supplement for 12 weeks in circulating endotoxin levels and cardiometabolic profiles of medication naïve T2DM patients: a randomized clinical trial. *J Transl Med* 2017; 15: 249.
- 11 Sabico S, Al-Mashharawi A, Al-Daghri NM, et al. Effects of a 6-month multi-strain probiotics supplementation in endotoxemic, inflammatory and cardiometabolic status of T2DM patients: A randomized, double-blind, placebo-controlled trial. *Clin Nutr* 2018; published online Aug 17. DOI:10.1016/j.clnu.2018.08.009.
- 12 Szulińska M, Łoniewski I, van Hemert S, Sobieska M, Bogdański P. Dose-Dependent Effects of Multispecies Probiotic Supplementation on the Lipopolysaccharide (LPS) Level and Cardiometabolic Profile in Obese Postmenopausal Women: A 12-Week Randomized Clinical Trial. *Nutrients* 2018; 10: 773.
- 13 Szulińska M, Łoniewski I, Skrypnik K, et al. Multispecies Probiotic Supplementation Favorably Affects Vascular Function and Reduces Arterial Stiffness in Obese Postmenopausal Women—A 12-Week Placebo-Controlled and Randomized Clinical Study. *Nutrients* 2018; 10: 1672.
- 14 Horvath A, Leber B, Feldbacher N, et al. Effects of a multispecies synbiotic on glucose metabolism, lipid marker, gut microbiome composition, gut permeability, and quality of life in diabetes: a randomized, double-blind, placebo-controlled pilot study. *Eur J Nutr* 2019; published online Nov 15. DOI:10.1007/s00394-019-02135-w.
- 15 Van Hemert S. Design of a multispecies probiotic product improving the intestinal Barrier. San Antonio, USA, 2012.
- 16 Hemert SV, Ormel G. Influence of the Multispecies Probiotic Ecologic® Barrier on Parameters of Intestinal Barrier Function. *Food Nutr Sci* 2014; 05: 1739.